

GEOLOGY DE
STANFORD UNI

CONTRIBUTIONS
FROM THE
CUSHMAN LABORATORY
FOR
FORAMINIFERAL RESEARCH

INDEX TO VOLUMES 6-10

PRICE \$1.00 POSTPAID

SHARON, MASSACHUSETTS, U. S. A.

CONTRIBUTIONS FROM THE CUSHMAN
LABORATORY FOR FORAMINIFERAL RESEARCH

INDEX TO VOLUMES 6—10 INCLUSIVE

Owing to the demand of those who wish to bind these Contributions, the following Index has been prepared.

All new genera and species appear in bold-faced type, others in italic type. An abbreviated form gives the geologic formation and general locality with each species.

Papers are grouped both by geographic and geologic headings as well as by subjects.

In order to conserve space, the following abbreviated form has been used. Volume number with the part number following in parentheses followed by a colon. After the colon comes the page number, then the plate number in bold-faced type, and the figure number in italics. It is hoped that this will give ready reference to the subject matter of these volumes.

- Actinosiphon*, 6(4) : 92; *semmesi* (Eoc.-Mexico), 6(4) : 92.
[Alabama]. An Eocene Foraminiferal Fauna of Wilcox Age from Alabama, 8(3) : 51. New Foraminifera from the Upper Jackson Eocene of the Southeastern Coastal Plain Region of the United States, 9(1) : 1.
- Ammobaculites alexanderi* (Cret.-Texas), 9(3) : 51, 5, 5; *arenata* (Cret.-Ark.), 9(3) : 50, 3, 4; *colombiana* (Cret.-Colombia), 6(3) : 68, 9, 4; *goodlandensis* (L. Cret.-Texas), 6(1) : 8, 2, 7, 8; *stephensoni* (Cret.-Texas), 9(3) : 49, 5, 2; *subcretacea* (L. Cret.-Texas), 6(1) : 6, 2, 9, 10; *texana* (Cret.-Texas), 9(3) : 50, 5, 3.
- Ammodiscus cretacea* 10(2) : 45; *incertus* (Rec.-Calif.), 6(3) : 51.
- Ammomassilina*, 9(2) : 32; *alveoliniformis*, 9(2) : 32, 3, 5.
- Ammospirata*, 9(2) : 32; *mexicana*, 9(2) : 32, 3, 4.

- Angulogerina albatrossi* (Rec.-Pac.), 8(2): 45, 6, 11, 12; var. *ornata* (Rec.-Pac.), 8(2): 45, 6, 13, 14; *angulosa* (Rec.-Calif.), 6(3): 60, 8, 7; *carinata* (Rec.-Pac.), 8(2): 44, 6, 7, 8; var. *bradyana* (Rec.-Pac.), 8(2): 45, 6, 9, 10; *fijiensis* (Plioc.-Fiji), 7(2): 31, 4, 11; *ocalana* (Eoc.-Ga.), 9(1): 14, 1, 30; *occidentalis* (Rec.-Pac.), 8(2): 46, 6, 15, 16.
- Angulogerinas from the Eastern Pacific, Some Recent, 8(2): 44.
- Anomalina affinis* (Eoc.-Calif.), 10(3): 64, 9, 2; *barrowi* (Eoc.-Texas), 7(3): 57, 7, 10; *clementiana* (Cret.-Antigua), 7(2): 46, 6, 10; *coalingensis*, 10(3): 63, 9, 3; *hughesi* (Mioc.-Calif.), 10(1): 23, 3, 13; *jacksonensis*, var. *limbosa* (Eoc.-Texas), 7(3): 58, 7, 11; *lorneiana* (Cret.-Antigua), 7(2): 45, 6, 9; *umbonifera* (Eoc.-Ala.), 8(3): 72, 9, 11.
- Annulocibicides*, 8(1): 1; *projectus* (Mioc.-Fla.), 8(1): 2, 1, 1.
- Annulopatulina*, 6(4): 87; *annularis* (Rec.-Australia), 6(4): 87. [Antigua]. Cretaceous Foraminifera from Antigua, B. W. I., 7(2): 33.
- Apterrinella*, 6(4): 82; *grahamensis* (Penn.-Okla., Texas), 6(4): 82, 11, 4-6.
- Arenobulimina brevicona*, 10(2): 35, 6, 7; 36; *conoidea*, 10(2): 36, 6, 9; *d'Orbigny*, 10(2): 30, 5, 14, 15; *frons*, 10(2): 31, 6, 2; *obesa*, 10(2): 31, 5, 20; *obliqua* (Cret.-Antigua), 7(2): 36, 5, 5; 10(2): 28, 5, 5, 6; *preslii*, 10(2): 29, 5, 12, 13; *puschi*, 10(2): 30, 5, 18; *sabulosa*, 10(2): 32, 6, 6; *subsphaerica*, 10(2): 30, 5, 16; *truncata*, 10(2): 29, 5, 8, 9.
- [Arkansas]. New American Cretaceous Foraminifera, 9(3): 49. Notes on Upper Cretaceous Species of Vaginulina, Flabellina and Frondicularia from Texas and Arkansas, 6(2): 25. *Rectogumbelina*, A New Genus from the Cretaceous, 8(1): 4.
- Articulina conico-articulata* (Rec.-Medit.), 7(3): 69, 9, 6; *terquemi* (Eoc.-Miss.), 9(1): 3, 1, 7.
- Asterodiscocyclina*, 6(4): 93; *stewarti* (Eoc.-Peru), 6(4): 93.
- Ataxophragmium rimosum*, 10(2): 32, 6, 3-5; *variabile* (Cret.-Antigua), 7(2): 36, 5, 6; 10(2): 28, 5, 2-4.
- Baculogypsinoides*, 6(4): 91; *spinosus* (Rec.-Indo-Pac.), 6(4): 91, 12, 10.
- Baggina californica* (Mioc.-Calif.), 7(1): 14, 2, 11.
- Bathysiphon alexanderi* (Cret.-Texas), 9(3): 49, 5, 1; *filiformis* (Mioc.-Calif.), 8(2): 32, 5, 1; *sanctaecrucis* (Mioc.-Calif.), 10(1): 1, 1, 1, 2; *taurinensis* (Cret.-Texas), 7(4): 88, 11, 12, 13.
- Bigenerina wintoni* (L. Cret.-Texas), 6(1): 9, 2, 15.

Bolivina advena, var. *striatella* (Rec.-Calif.), 6(3): 58, 8, 6; (Mioc.-Calif.), 7(1): 9; *argentea* (Rec.-Calif.), 6(3): 57, 8, 3; *barbarana* (Mioc.-Calif.), 10(1): 11, 2, 5; *capitata* (Rec.-Pac.), 9(4): 80, 8, 12; *costata*, var. *bicostata* (Rec.-Calif.), 6(3): 57; var. *interjuncta* (Rec.-Calif.), 6(3): 57; *explicata* (Cret.-Venezuela), 6(3): 66, 9, 8, 9; *floridana* (Mioc.-Calif.), 7(1): 9, 2, 2; *girardensis* (Mioc.-Calif.), 10(1): 17, 3, 7; *globulosa* (Rec.-Pac.), 9(4): 80, 8, 9; *goudkoffi* (Mioc.-Calif.), 10(1): 22, 4, 4, 5; *granti* (Mioc.-Calif.), 10(1): 21, 4, 2, 3; *hootsi* (Mioc.-Calif.), 10(1): 18, 4, 1; *hughesi*, var. *parva* (Mioc.-Calif.), 10(1): 25, 4, 10; *marginata* (Mioc.-Calif.), 7(1): 9, 2, 1; var. *adelaidana* (Mioc.-Calif.), 10(1): 10, 2, 1, 2; var. *gracilis* (Mioc.-Calif.), 10(1): 10, 2, 3; *modeloensis* (Mioc.-Calif.), 10(1): 10, 2, 4; *oceanica* (Rec.-Pac.), 9(4): 81, 8, 10; *pseudo-beyrichii* (Rec.-Calif.), 6(3): 58, 8, 5; *pseudopygmaea* (Rec.-Pac.), 9(4): 79, 8, 8; *pygmaea* (Rec.-Calif.), 6(3): 57, 8, 4; *seminuda* (Rec.-Calif.), 6(3): 57; *subadvena* (Rec.-Calif.), 6(3): 57; var. *spissa* (Rec.-Calif.), 6(3): 58; *vadescens* (Rec.-Pac.), 9(4): 81, 8, 11.

Bolivinita eleyi (Cret.-Antigua), 7(2): 39, 5, 8; *selmensis* (Cret.-Tenn.), 9(3): 58, 7, 3, 4; *subangularis*, var. *lineata* (Rec.-Pac.), 9(4): 77, 8, 1, 2.

Bolivinoides velascoensis, 8(4): 94.

Bonairea Pijpers, 1933, *Ruttenia*, A New Name for, 9(2): 30.

Bulimina, Notes on Some of the Earlier Species Originally Described as, 10(2): 27.

Bulimina acuta, 10(2): 31, 5, 21; *amphicona*, 10(2): 36; *brevis* (Cret.-Antigua), 7(2): 40, 5, 9; 10(2): 28; *capitata* (Eoc.-Calif.), 10(3): 61, 8, 10; *carnerosensis* (Mioc.-Calif.), 10(1): 5, 1, 12; var. *mahoneyi* (Mioc.-Calif.), 10(1): 5, 1, 13; *cenomana*, 10(2): 36; *cooperensis* (Eoc.-So. Car.), 9(1): 12, 1, 32; *del-reyensis* (Mioc.-Calif.), 10(1): 25, 4, 8; *fijiensis* (Rec.-Pac.), 9(4): 79, 8, 7; *imbricata* (Cret.-Antigua), 7(2): 42, 5, 10; *inflata* (Mioc.-Calif.), 7(1): 9; *murchisoniana*, 10(2): 29, 5, 7; *ovata* (Rec.-Calif.), 6(3): 56, 7, 22; (Eoc.-Ala.), 8(3): 67, 9, 1, 2; *ovula* (Rec.-Calif.), 6(3): 56, 7, 21; *ovulum*, 10(2): 29, 5, 10, 11; *pulchella* (Rec.-Calif.), 6(3): 56, 7, 19; *puschi* (Cret.-Antigua), 7(2): 42, 5, 11; *sarthacensis*, 10(2): 36; *schwageri* (Eoc.-Calif.), 10(3): 62, 8, 11; *tortilis*, 10(2): 31, 6, 1; *trigona*, 10(2): 36; *tumida*, 10(2): 29; *uvigerinaformis* (Mioc.-Calif.), 10(1): 5, 1, 14.

Buliminella colonensis (Cret.-Venezuela), 6(3): 65, 9, 6, 7; *d'Orbigny* (Cret.-Venezuela), 6(3): 65, 9, 3; cf. *elegantissima*

- (Eoc.-Ala.), 8(3): 67, 8, 20, 21; *henryana* (Mioc.-Calif.), 10(1): 4, 1, 11; *imbricata*, 10(2): 31, 5, 19; *milletti* (Rec.-Pac.), 9(4): 78, 8, 5, 6; *obtusa*, 10(2): 28, 5, 1; *subfusiformis* (Rec.-Calif.), 6(3): 56, 7, 20; (Mioc.-Calif.), 7(1): 8, 1, 29.
- Calcitornella*, 6(4): 81; *elongata* (Penn.-Texas), 6(4): 81, 11, 7. *Calcivertella*, 6(4): 82; *adherens* (Penn.-Texas), 6(4): 82, 11, 3. [California]. A New *Spiroplectoides* from the Cretaceous of California, 10(3): 70. Additional New Foraminifera from the Miocene of California, 10(1): 24. Eocene Foraminifera of the Poway Conglomerate of California, 10(3): 51. Miocene Foraminifera from the Temblor of the East Side of the San Joaquin Valley, California, 7(1): 1. New and Unrecorded Foraminifera from the California Miocene, 10(1): 1. Notes on Some Arenaceous Foraminifera from the Temblor Formation of California, 8(2): 29. Some Recent Foraminifera from off San Pedro, California, 6(3): 49.
- Cancris baggi* (Mioc.-Calif.), 10(1): 15, 3, 2.
- Cassidulina angulosa* (Rec.-Pac.), 9(4): 93, 10, 6; *barbarana* (Mioc.-Calif.), 10(1): 16, 3, 5; *costatula* (Rec.-Pac.), 9(4): 94, 10, 7; *delicata* (Rec.-Calif.), 6(3): 61, 8, 16; *laevigata* (Rec.-Calif.), 6(3): 61; var. *carinata* (Mioc.-Calif.), 7(1): 14, 2, 14; *limbata* (Rec.-Calif.), 6(3): 61; *margareta* (Mioc.-Calif.), 7(1): 15, 2, 13; *minuta* (Rec.-Pac.), 9(4): 92, 10, 3; *modeloensis* (Mioc.-Calif.), 10(1): 23, 3, 12; *monicana* (Mioc.-Calif.), 10(1): 16, 3, 4; *patula* (Rec.-Pac.), 9(4): 92, 10, 2; *rariocula* (Rec.-Pac.), 9(4): 93, 10, 4; *subtumida* (Rec.-Pac.), 9(4): 93, 10, 5; *translucens* (Rec.-Calif.), 6(3): 60.
- Chilostomella grandis* (Rec.-Calif.), 6(3): 61, 8, 14; *oolina* (Rec.-Calif.), 6(3): 61, 8, 15.
- Chrysalogonium breviloculum* (Mioc.-Trinidad), 10(3): 74, 10, 13; *elongatum* (Mioc.-Trinidad), 10(3): 73, 10, 10, 11; *lanceolum* (Mioc.-Trinidad), 10(3): 75, 10, 16; *longicostatum* (Mioc.-Trinidad), 10(3): 74, 10, 12.
- Cibicides americanus* (Mioc.-Calif.), 7(1): 15, 3, 1; *cooperensis* (Eoc.-So. Car.), 9(1): 20, 2, 11; *floridanus* (Mioc.-Calif.), 7(1): 16, 3, 2; *gallowayi* (Rec.-Calif.), 6(3): 62; *martinezensis* (Eoc.-Calif.), 10(3): 64; *praecursorius* (Eoc.-Ala.), 8(3): 72, 9, 14; *pseudowuellerstorfi* (Eoc.-Calif.), 10(3): 64, 9, 4; *sassei* (Eoc.-Calif.), 10(3): 64, 9, 5; *semiplectus* (Eoc.-Ala.), 8(3): 72, 9, 12, 13 (?); *ungeriana* (Rec.-Calif.), 6(3): 62; *yazooensis* (Eoc.-Miss.), 7(3): 59, 7, 12.
- Clavulina* from the Cretaceous of Texas, A New Species of, 9(1): 21.

Clavulina arenata (Cret.-Ark.), 9(3) : 54, 6, 5; *communis* (Mioc.-Calif.), 8(2) : 38, 5, 15, 16; *disjuncta* (Cret.-Texas), 9(1) : 22; *flintiana* (Mioc.-Calif.), 8(2) : 39, 5, 17; *flintii* (Rec.-Calif.), 6(3) : 52, 7, 7-9; *humilis*, var. *mexicana* (Mioc.-Calif.), 8(2) : 39, 5, 18, 19.

[Colombia]. Notes on Some Foraminifera from Venezuela and Colombia, 6(3) : 64.

Colonammia, 6(4) : 75; *verruca* (Pal.-Amer.), 6(4) : 75, 10, 4.

Conicospirillina coronata (Rec.-Pac.), 9(4) : 87, 9, 1; *semi-involuta* (Rec.-Pac.), 9(4) : 86, 8, 19; *trochoidea* (Rec.-Pac.), 9(4) : 87, 8, 20.

"*Cornuspira cretacea* Reuss," The Generic Position of, 10(2) : 44.

Cornuspira foliacea (Rec.-Calif.), 6(3) : 52, 7, 12.

[Cretaceous]. A New Species of *Clavulina* from the Cretaceous of Texas, 9(1) : 21. A New *Spiroplectoides* from the Cretaceous of California, 10(3) : 70. Cretaceous Foraminifera from Antigua, B. W. I., 7(2) : 33. *Hastigerinella* and Other Interesting Foraminifera from the Upper Cretaceous of Texas, 7(4) : 83. New American Cretaceous Foraminifera, 9(3) : 49. Notes on Some Foraminifera from Venezuela and Colombia, 6(3) : 64. Notes on Some of the Earlier Species Originally Described as *Bulimina*, 10(2) : 27. Notes on Upper Cretaceous Species of *Vaginulina*, *Flabellina* and *Fronicularia* from Texas and Arkansas, 6(2) : 25. *Rectogumbelina*, A New Genus from the Cretaceous, 8(1) : 4. Some *Vaginulinas* and Other Foraminifera from the Lower Cretaceous of Texas, 6(1) : 1. *Textularia* and Related Forms from the Cretaceous, 8(4) : 86. The Development of *Hantkenina* in the Cretaceous, with a Description of a New Species, 6(2) : 39. The Generic Position of "*Cornuspira cretacea* Reuss," 10(2) : 44. The Relationships of *Textulariella* and Description of a New Species, 8(4) : 97. Two New Navarro Foraminifera from Texas, 8(4) : 98.

Cyclammia incisa (Mioc.-Calif.), 8(2) : 32, 5, 2.

Cycloloculina miocenica (Mioc.-Fla.), 8(1) : 3, 1, 3, 4.

Deckerella, 6(4) : 78; *clavata* (Penn.-Texas), 6(4) : 78, 10, 11.

Dentalina barnesi (Mioc.-Calif.), 10(1) : 17, 3, 6; *capitata* (Eoc.-Calif.), 10(3) : 56, 7, 20; *colei* (Eoc.-Calif.), 10(3) : 54, 7, 10-12; *communis* (Mioc.-Calif.), 7(1) : 3, 1, 8; (Rec.-Medit.), 7(3) : 68, 8, 22, 23; (Eoc.-Ala.), 8(3) : 55, 7, 12, 13; *consobrina* (Eoc.-Calif.), 10(3) : 55, 7, 13-15; *cooperensis* (Eoc.-So. Car.), 9(1) : 8, 1, 17; *halkyardi* (Eoc.-So. Car.), 9(1) : 9, 1, 20; *hantkeni* (Eoc.-Ala.), 9(1) : 9, 1, 18, 19; *jacksonensis* (Eoc.-Ala.), 8(3) : 55, 7, 10, 11; *obliquisuturata* (Eoc.-Calif.), 10(3) : 54, 7, 22-25;

quadrulata (Mioc.-Calif.), 7(1) : 3, 1, 9-11; *substrigata* (Eoc.-Calif.), 10(3) : 55, 7, 17-19; *vagina* (Eoc.-Calif.), 10(3) : 55, 7, 16; *vertebralis* (Rec.-Medit.), 7(3) : 66, 8, 20, 21; sp(?) (Eoc.-Ala.), 8(3) : 56, 7, 17.

Dentostomina, A New Genus of the Miliolidae, 9(2) : 31.

Dentostomina, 9(2) : 31; *bermudiana* (Rec.-Atl.), 9(2) : 31, 3, 6.

Discorbis alabamensis (Eoc.-Ala.), 9(1) : 16, 2, 3; *alveata* (Eoc.-Miss.), 9(1) : 16, 2, 4; *assulata* (Eoc.-Ga.), 9(1) : 15, 2, 2; *bullata* (Eoc.-Ala.), 9(1) : 16, 2, 6; *crustata* (Rec.-Pac.), 9(4) : 88, 9, 4; *farishi* (Eoc.-La.), 8(2) : 43, 6, 6; *frustata* (Rec.-Pac.), 9(4) : 88, 9, 2; *globulo-spinosa* (Eoc.-Miss.), 9(1) : 14, 2, 1; *hemisphaerica* (Eoc.-Miss.), 7(3) : 59, 7, 14; *isabelleana* (Rec.-Calif.), 6(3) : 60, 8, 8; *micens* (Rec.-Pac.), 9(4) : 89, 9, 5; *ocalana* (Eoc.-Fla.), 9(1) : 15, 2, 5; *opima* (Rec.-Pac.), 9(4) : 88, 9, 3.

Dorothia californica (Mioc.-Calif.), 10(1) : 1, 1, 18; *conulus*, 8(4) : 89; *glabrata* (Cret.-Texas), 9(3) : 56, 6, 10; *glabrella* (Cret.-Texas), 9(3) : 56, 6, 9; *pontoni* (Cret.-Miss.), 9(3) : 55, 6, 8; *pupa*, 8(4) : 91.

Dyocibicides, 6(4) : 91; *biserialis* (Rec.-Pac.), 6(4) : 91, 12, 12.

Earlandia, 6(4) : 75; *perparva* (Penn.-Texas), 6(4) : 75.

Eggerella, 9(2) : 33; *bradyi*, 9(2) : 33, 4, 1.

Ehrenbergina albatrossi (Rec.-Pac.), 9(4) : 94, 10, 8; *reticulata* (Rec.-Pac.), 9(4) : 94, 10, 9.

Ellipsolagena fijiensis (Plioc.-Fiji), 7(2) : 32, 4, 6.

Ellipsonodosaria dentaliniformis (Mioc.-Trinidad), 10(3) : 73, 10, 9; *mappa* (Mioc.-Trinidad), 10(3) : 73, 10, 8; *nuttalli* (Mioc.-Trinidad), 10(3) : 72, 10, 6; var. *gracillima* (Mioc.-Trinidad), 10(3) : 72, 10, 7.

Elphidium advenum, var. *margaritaceum* (Pleistoc.-Md.), 6(4) : 97; *discoideale* (Pleistoc.-Md.), 6(4) : 97, 13, 10; *eocenicum* (Eoc.-Texas), 7(3) : 53, 7, 6; *incertum* (Pleistoc.-Md.), 6(4) : 96, 13, 6, 7; var. *clavatum* (Pleistoc.-Md.), 6(4) : 96, 13, 8, 9; *schlencki* (Eoc.-Calif.), 10(3) : 60, 8, 8; *smithi* (Eoc.-Calif.), 10(3) : 61, 8, 7; *spinatum* (Rec.-Calif.), 6(3) : 56; *subsphaericum* (Mioc.-Venezuela), 6(3) : 64, 9, 1, 2.

Endothyra baileyi (Hall), The Species, 7(4) : 90.

Endothyra baileyi (Pal.-Ind.), 7(4) : 90, 11, 16-28; 12, 3-8.

Endothyranella, 6(4) : 76; *powersi* (Penn.-Okla., Texas), 6(4) : 76, 10, 15.

Entosolenia lucida (Pleistoc.-Md.), 6(4) : 98, 13, 11, 12.

[Eocene]. A New Genus of the Foraminifera, Gunteria, from the Middle Eocene of Florida, 9(2) : 25. Additional New Eocene

Foraminifera, 8(2): 40. An Eocene Foraminiferal Fauna of Wilcox Age from Alabama, 8(3): 51. Eocene Foraminifera of the Poway Conglomerate of California, 10(3): 51. Fossil Species of Hastigerinella, 6(1): 17. New Foraminifera from the Upper Jackson Eocene of the Southeastern Coastal Plain Region of the United States, 9(1): 1. New Species of Triloculina from the Claiborne of Louisiana, 10(3): 65. Post-Cretaceous Occurrence of Gumbelina with a Description of a New Species, 9(3): 64. Some New Eocene Foraminifera from Jamaica, 7(4): 75. Some New Tertiary Foraminifera from Texas, 7(3): 51. The Range of Sigmoidella plummerae Cushman and Ozawa. A Correction, 6(4): 101. Three New Upper Eocene Foraminifera, 7(3): 58. Two New Texas Foraminifera, 9(4): 95.

Eoguttulina, 6(4): 86; *anglica* (Cret.-England), 6(4): 86, 12, 3. *Eouvigerina aculeata* (Cret.-Texas), 9(3): 62, 7, 8; *austinana* (Cret.-Texas), 9(3): 61, 7, 5; *plummerae* (Cret.-Texas), 9(3): 62, 7, 6, 7.

Epistomina ramonensis (Mioc.-Calif.), 10(1): 15, 3, 1.

Eponides exigua (Rec.-Calif.), 6(3): 60, 8, 11; *fijiana* (Rec.-Pac.), 9(4): 89, 9, 6; *frigida*, var. *calida* (Pleistoc.-Md.), 6(4): 98, 13, 13; *guayabalensis*, var. *yeguaensis* (Eoc.-Calif.), 10(3): 62, 9, 1; *jamaicensis* (Eoc.-Jamaica), 7(4): 77, 10, 4; *keenani* (Mioc.-Calif.), 10(1): 14, 3, 10, 11; *lotus* (Eoc.-Ala.), 8(3): 71, 9, 8; *mansfieldi* (Mioc.-Calif.), 7(1): 12, 2, 10; *minima* (Eoc.-So. Car.), 9(1): 17, 2, 8; *ocalana* (Eoc.-Ala.), 9(1): 17, 2, 7; *ornata* (Rec.-Calif.), 6(3): 60, 8, 12; *rosaformis* (Mioc.-Calif.), 10(1): 14, 2, 18; *tenera* (Mioc.-Calif.), 10(1): 15, 2, 19; *terebra* (Rec.-Pac.), 9(4): 89, 10, 1; *vicksburgensis* (Olig.-Texas), 7(3): 56, 7, 8.

Flabellina and Frondicularia from Texas and Arkansas, Notes on Upper Cretaceous Species of Vaginulina, 6(2): 25.

Flabellina crassa (Mioc.-Calif.), 10(1): 3, 1, 6; *harpa* (Rec.-Medit.), 7(3): 70, 9, 15; *interpunctata* (Cret.-Texas, etc.), 6(2): 30, 4, 16, 17; *reticulata* (Cret.-Texas, Ark.), 6(2): 32, 4, 18, 19; *rugosa* (Cret.-Texas, etc.), 6(2): 32, 4, 15; (Cret.-Antigua), 7(2): 38, 5, 3.

Flabellinella Schubert, Some Notes on the Genus, 7(1): 16.

Flabellinella zitteliana (Cret.-Germany), 7(1): 16, 3, 3-5.

[Florida]. A New Genus of the Foraminifera, Gunteria, from the Middle Eocene of Florida, 9(2): 25. A New Plectofrondicularia from Florida, 7(3): 60. A New Virgulina from the Miocene of Florida, 7(2): 32. New Foraminifera from the Upper

- Jackson Eocene of the Southeastern Coastal Plain Region of the United States, 9(1): 1. Some Interesting New Foraminifera from the Miocene of Florida, 8(1): 1.
- Franklina*, 6(4): 77; *goodlandensis* (Cret.-Europe, Amer.), 6(4): 77, 10, 12.
- Fronicularia* from Texas and Arkansas, Notes on Upper Cretaceous Species of *Vaginulina*, *Flabellina*, and, 6(2): 25.
- Fronicularia angusta* (Cret.-Ark.), 6(2): 38, 5, 21; *archiaciana* (Cret.-Tex., Ark.), 6(2): 37, 5, 9-12; *clarki* (Cret.-Texas), 6(2): 34, 5, 1, 2; *complanata* (Rec.-Medit.), 7(3): 70, 9, 14; *cordai* (Cret.-Texas), 6(2): 34, 5, 17; *decheni* (Cret.-Texas, Ark.), 6(2): 38, 5, 20; *goldfussi* (Cret.-Texas, Ark.), 6(2): 33, 5, 3; *gracilis* (Cret.-Texas), 6(2): 37, 5, 16; *lanceola* (Cret.-Texas), 6(2): 38, 5, 18, 19; *microdisca* (Cret.-Texas), 6(2): 33, 5, 4; *verneuilliana* (Cret.-Texas, Ark.), 6(2): 36, 5, 5, 6; var. *bidentata* (Cret.-Texas), 6(2): 37, 5, 13-15; var. *fossata* (Cret.-Texas), 6(2): 36, 5, 7, 8.
- Gaudryina atlantica* (Mioc.-Calif.), 8(2): 37, 5, 14; var. *asperita* (Mioc.-Calif.), 8(2): 37, 5, 13; *bentonensis*, 8(4): 96; *concinna*, 8(4): 90; *faujasi*, 8(4): 91; *foeda*, 8(4): 90; *io* (Cret.-Texas), 9(3): 51, 5, 6; *lobulata* (Mioc.-Calif.), 8(2): 38, 5, 12; *navarroana* (Cret.-Texas, Ark.), 8(4): 98, 11, 15, 16; *subquadrata* (Eoc.-So. Car.), 9(1): 2, 1, 1; *subrotundata* (Mioc.-Calif.), 8(2): 34, 5, 8, 9; *triangularis* (Rec.-Calif.), 6(3): 52, 7, 5; (Mioc.-Calif.), 7(1): 2, 1, 1; (Mioc.-Calif.), 8(2): 36, 5, 11; *trinitatensis* (Mioc.-Calif.), 8(2): 36, 5, 4, 5, 10.
- Gaudryinella capitosa* (Cret.-Texas), 9(3): 52, 5, 8; var. *serulata* (Cret.-Texas), 9(3): 52, 5, 9; *mollis* (Cret.-Miss.), 9(3): 53, 6, 6; *pseudoserrata* (Cret.-Texas), 8(4): 99, 11, 20, 21.
- Glandulina ozawai* Cushman, New Species, 7(4): 83.
- Glandulina abbreviata* (Eoc.-Ala.), 8(3): 64, 8, 9; *laevigata* (Mioc.-Calif.), 7(1): 7, 1, 25; (Eoc.-Ala.), 8(3): 64, 8, 10; var. *ovata* (Eoc.-Calif.), 10(3): 60; *ozawai* (Olig.-Germany), 7(4): 83.
- Globigerina conglomerata* (Rec.-Calif.), 6(3): 61; *cretacea* (Cret.-Antigua), 7(2): 44, 6, 6; 10(3): 63, 8, 13; *elevata* (Cret.-Antigua), 7(2): 44, 6, 7.
- Globigerinella aspera* (Cret.-Antigua), 7(2): 44, 6, 5.
- Globobulimina pacifica* (Rec.-Calif.), 6(3): 57; (Mioc.-Calif.), 7(1): 9, 1, 30.
- Globorotalia micheliniana* (Cret.-Antigua), 7(2): 45, 6, 8; *truncatulinoides* (Rec.-Calif.), 6(3): 62; *wilcoxensis* (Eoc.-Ala.), 8(3): 71, 9, 10.

- Globulina gibba*, var. (Eoc.-Calif.), 10(3) : 59, 8, 4; *landesi* (Eoc.-Calif.), 10(3) : 59, 8, 1-3.
- Glyphostomella*, 6(4) : 86; *triloculina* (Penn.-Texas), 6(4) : 86, 12, 1.
- Goësella**, 9(2) : 34; *rotundata*, 9(2) : 34, 4, 2; *rugulosa* (Cret.-Miss.), 9(3) : 55, 6, 7.
- Guembeina globulosa* (Cret.-Antigua), 7(2) : 39, 5, 7.
- Gümbelina with a Description of a New Species, Post-Cretaceous Occurrence of, 9(3) : 64.
- Gümbelina excolata*, 8(4) : 94; *globifera*, 8(4) : 91; *globulosa*, 8(4) : 90; **goodwini** (Eoc.-Trinidad), 9(3) : 69, 7, 15, 16; *striata*, 10(3) : 61, 8, 9; *wilcoxensis* (Eoc.-Ala.), 8(3) : 66, 8, 16, 17.
- Gümbelitria (?) from the Pacific, A Recent, 10(4) : 105.
- Gümbelitria**, 9(2) : 37; **cretacea** (Cret.-Texas), 9(2) : 37, 4, 12; (?) **vivans** (Rec.-Pac.), 10(4) : 105, 13, 9, 10.
- Gunteria, from the Middle Eocene of Florida, A New Genus of the Foraminifera, 9(2) : 25.
- Gunteria**, 9(2) : 25; **floridana** (Eoc.-Fla.), 9(2) : 26, 3, 1-3.
- Guttulina hantkeni* (Eoc.-Calif.), 10(3) : 60, 8, 5; *problema* (Eoc.-Ala.), 8(3) : 61, 8, 3, 4; *quinquecosta* (Rec.-Calif.), 6(3) : 53, 7, 15; **wilcoxensis** (Eoc.-Ala.), 8(3) : 60, 8, 1, 2.
- Gyroidina soldanii* (Mioc.-Calif.), 7(1) : 11, 2, 9; var. *octocamerata* (Eoc.-Calif.), 10(3) : 62; *umbilicata* (Cret.-Antigua), 7(2) : 43, 6, 3.
- Hantkenina in the Cretaceous with a Description of a New Species, The Development of, 6(2) : 39.
- Hantkenina cenomana* (Cret.-Germany), 6(2) : 40, 6, 1-3; **multi-spinata** (Cret.-Canada), 6(2) : 40, 6, 4-6; (Cret.-Texas), 7(4) : 88, 11, 10, 11.
- Haplophragmoides advena* (Rec.-Calif.), 6(3) : 51, 7, 4; *planissima* (Rec.-Calif.), 6(3) : 51, 7, 3.
- Haplostiche jamaicensis* (Eoc.-Jamaica), 7(4) : 75, 10, 1.
- Hastigerinella and Other Interesting Foraminifera from the Upper Cretaceous of Texas, 7(4) : 83. Fossil Species of *Hastigerinella*, 6(1) : 17.
- Hastigerinella alexanderi* (Cret.-Texas), 7(4) : 87, 11, 6-9; *eocanica* (Eoc.-Trinidad), 6(1) : 18, 3, 6, 7; **jarvisi** (Eoc.-Trinidad), 6(1) : 18, 3, 8-11; **moremani** (Cret.-Texas), 7(4) : 86, 11, 1-3; **watersi** (Cret.-Texas), 7(4) : 86, 11, 4, 5.
- Heterostomella austinana* (Cret.-Texas), 9(3) : 53, 6, 1-3; *boynensis* (Cret.-Canada), 9(3) : 54; *cuneata* (Cret.-Ala.),

- 9(3) : 54; var. *curvata* (Cret.-Miss.), 9(3) : 54, 6, 4; *foveolata*, 9(3) : 54.
- Hopkinsina pacifica* (Rec.-Pac.), 9(4) : 86, 8, 16.
- Hyperamminoides*, 6(4) : 75; *elegans* (Penn.-Amer.), 6(4) : 75, 10, 8, 9.
- [Jamaica.] Some New Eocene Foraminifera from Jamaica, 7(4) : 75.
- Karrerella*, 9(2) : 34; *siphonella*, 9(2) : 34, 4, 3, 4.
- Kyphopyxa*, 6(4) : 85; *christneri* (Cret.-Texas, etc.), 6(2) : 33, 4, 20; (Cret.-Venezuela), 6(3) : 65, 9, 5; (Cret.-Texas), 6(4) : 85, 12, 2.
- Lagena acuticosta* (Eoc.-Ala.), 8(3) : 59, 7, 20; *basi-striatula* (Plioc.-Fiji), 7(2) : 27, 4, 5; *clavata* (Eoc.-Ala.), 8(3) : 60, 7, 23; *costata* (Mioc.-Calif.), 7(1) : 6, 1, 21; (Eoc.-Ala.), 8(3) : 59, 7, 21; *elongata* (Rec.-Calif.), 6(3) : 53, 7, 14; *gracilis* (Rec.-Calif.), 6(3) : 53; *hexagona*, var. *scalariformis* (Rec.-Calif.), 6(3) : 53; (Mioc.-Calif.), 7(1) : 7, 1, 24; *hispida* (Rec.-Calif.), 6(3) : 53, 7, 13; *lamellata* (Rec.-Calif.), 6(3) : 53; *mucronulata* (Eoc.-Ala.), 8(3) : 59, 7, 18; *orbignyana*, var. (Eoc.-Ala.), 8(3) : 60, 7, 24; var. *semiconcentrica* (Eoc.-No. Car.), 9(1) : 10, 1, 22; *perlucida* (Mioc.-Calif.), 7(1) : 6, 1, 22; *schwageriana* (Plioc.-Fiji), 7(2) : 26, 4, 2; *semistriata* (Mioc.-Calif.), 7(1) : 7, 1, 23; *spino-alata* (Plioc.-Fiji), 7(2) : 27, 4, 4; *spiro-striolata* (Plioc.-Fiji), 7(2) : 27, 4, 3; *sulcata* (Mioc.-Calif.), 7(1) : 6, 1, 20; var. *apiculata* (Rec.-Calif.), 6(3) : 53; sp(?) (Eoc.-Ala.), 8(3) : 60, 7, 22.
- Lagenammina sphaerica* (Pal.-Okla.), 6(2) : 6, 10; *stillia* (Pal.-Okla.), 6(2) : 6, 9.
- Lamarckina jacksonensis* (Eoc.-Ala.), 9(1) : 20, 2, 10; *wilcoxensis* (Eoc.-Ala.), 8(3) : 70, 9, 4.
- Lenticulina* (?) *convergens* (Eoc.-Calif.), 10(3) : 54, 7, 7; *rotulata* (Cret.-Antigua), 7(2) : 37, 5, 1.
- Liebusella*, 9(2) : 36; *soldanii*, 9(2) : 36, 4, 9-11.
- Lingulina wilcoxensis* (Eoc.-Ala.), 8(3) : 58, 7, 14.
- Listerella*, 9(2) : 36; *primaeva*, 9(2) : 36, 4, 5.
- [Louisiana]. Additional New Eocene Foraminifera, 8(2) : 40.
New Species of Triloculina from the Claiborne of Louisiana, 10(3) : 65.
- Loxostoma rostrum* (Rec.-Pac.), 9(4) : 82, 8, 13.
- Loxostomum wilcoxensis* (Eoc.-Ala.), 8(3) : 67, 9, 3.
- Marginulina cooperensis* (Eoc.-So. Car.), 9(1) : 8, 1, 14, 15; *cooperi* (Eoc.-So. Car.), 9(1) : 5, 1, 13; *costatus* (Rec.-Medit.),

- 7(3) : 64, 8, 2-5; *dubia* (Mioc.-Calif.), 7(1) : 3, 1, 6; (Eoc.-Ala.), 8(3) : 53, 7, 6; *eximia* (Eoc.-Ala.), 8(3) : 54, 7, 8; *murex* (Rec.-Medit.), 7(3) : 65, 8, 9-11; *pachygaster* (Eoc.-Ala.), 8(3) : 53, 7, 4, 5; *subbullata* (Mioc.-Calif.), 7(1) : 3, 1, 7; (Eoc.-Ala.), 8(3) : 53, 7, 7.
- Marssonella**, 9(2) : 36; *oxycona*, 9(2) : 36, 4, 13.
- Martinottiella**, 9(2) : 37; *communis*, 9(2) : 37, 4, 6-8.
- [Maryland]. Pleistocene Foraminifera from Maryland, 6(4) : 94.
- Massilina** *humblei* (Eoc.-Texas), 8(2) : 40, 6, 1; *jacksonensis* (Eoc.-Miss.), 9(1) : 2, 1, 4; var. *punctato-costata* (Eoc.-Miss.), 9(1) : 3, 1, 5, 6; *pratti* (Eoc.-Texas), 7(3) : 53, 7, 4.
- Miliammina**, 6(4) : 81; *oblonga*, var. *arenacea* (Rec.-Antarctic), 6(4) : 81, 10, 19-21.
- Miliola** *jacksonensis* (Eoc.-Miss.), 9(1) : 2, 1, 2, 3.
- Mimosina** *pacifica* (Rec.-Pac.), 9(4) : 77, 8, 3.
- [Miocene]. A New Plectofrondicularia from Florida, 7(3) : 60. A New Virgulina from the Miocene of Florida, 7(2) : 32. Additional New Foraminifera from the Miocene of California, 10(1) : 24. Miocene Foraminifera from the Temblor of the East Side of the San Joaquin Valley, California, 7(1) : 1. New and Unrecorded Foraminifera from the California Miocene, 10(1) : 1. Notes on Some Arenaceous Foraminifera from the Temblor Formation of California, 8(2) : 29. Notes on Some Foraminifera from Venezuela and Colombia, 6(3) : 64. Some Interesting New Foraminifera from the Miocene of Florida, 8(1) : 1. Some Interesting New Uniserial Foraminifera from Trinidad, 10(3) : 71.
- [Mississippi]. New American Cretaceous Foraminifera, 9(3) : 49. New Foraminifera from the Upper Jackson Eocene of the Southeastern Coastal Plain Region of the United States, 9(1) : 1.
- Monalysidium** *lituus* (Rec.-Medit.), 7(3) : 72, 9, 22.
- Mooreinella**, 6(4) : 84; *biserialis* (Penn.-Texas), 6(4) : 84, 11, 11-13.
- Myogypsinoidea**, 6(4) : 93; *dehaartii* (Mioc.-Indo-Pac.), 6(4) : 93.
- Nautilus** (O.) *radicula* (Rec.-Medit.), 7(3) : 68, 9, 5; *vaginaeformis* (Rec.-Medit.), 7(3) : 68, 8, 14.
- Nodogenerina** *cooperensis* (Eoc.-So. Car.), 9(1) : 11, 1, 27; *milletti* (Rec.-Pac.), 9(4) : 86, 8, 17, 18.
- Nodosarella** *gracillima* (Cret.-Texas), 9(3) : 64, 7, 14; *pacifica* (Plioc.-Fiji), 7(2) : 31, 4, 12, 13; sp(?) (Cret.-Antigua), 7(2) : 42, 6, 4.

Nodosaria affinis (Cret.-Antigua), 7(2): 38, 5, 4; *anomala* (Mioc.-Calif.), 7(1): 4, 1, 12-14; *arundinea* (Mioc.-Calif.), 7(1): 6, 1, 17-19; *cookei* (Eoc.-Ala.), 9(1): 9, 1, 21; *ewaldi* (Eoc.-Calif.), 10(3): 56, 7, 21; *globifera* (Rec.-Medit.), 7(3): 68, 9, 1-4; *latejugata*, var. (Eoc.-Ala.), 8(3): 58, 7, 15, 16; var. *carolinensis* (Eoc.-So. Car.), 9(1): 5, 1, 16; *obliquata* (Rec.-Medit.), 7(3): 65, 8, 15-19; *parexilis* (Mioc.-Calif.), 7(1): 6, 1, 15; var. *sentifera* (Mioc.-Calif.), 7(1): 6, 1, 16; *scalaris* (Rec.-Medit.), 7(3): 65, 8, 12, 13; *tornata* (Mioc.-Calif.), 10(1): 3, 1, 7.

Nonion costiferum (Mioc.-Calif.), 7(1): 7, 1, 27; *galeata* (Plioc.-Fiji), 7(2): 28, 4, 7; *hantkeni*, var. *fayettei* (Eoc.-La.), 8(2): 41, 6, 3; *incisum* (Mioc.-Calif.), 7(1): 7, 1, 26; cf. *labradoricum* (Mioc.-Calif.), 10(1): 4, 1, 8; *laevis*, var. *marginatum* (Eoc.-Texas), 7(3): 52, 7, 5; *montereyanum* (Mioc.-Calif.), 10(1): 24, 4, 6; var. *carmeloensis* (Mioc.-Calif.), 10(1): 24, 4, 7; *pizarrensis* (Mioc.-Calif.), 10(1): 4, 1, 9; var. *basispinata* (Rec.-Calif.), 6(3): 54, 7, 18; var. *multicameratum* (Mioc.-Calif.), 10(1): 4, 1, 10; *planatum* (Eoc.-Calif.), 10(3): 60, 8, 6; *pompilioides* (Pleistoc.-Md.), 6(4): 95, 13, 4; *scaphum*, var. *inflatum* (Eoc.-La.), 8(2): 41, 6, 2; *sloanii* (Pleistoc.-Md.), 6(4): 96, 13, 5; *wilcoxensis* (Eoc.-Ala.), 8(3): 64, 8, 11.

Nonionella alabamensis (Eoc.-Ala.), 8(3): 65, 8, 15; *austinana* (Cret.-Texas), 9(3): 57, 7, 2; *clavata* (Plioc.-Fiji), 7(2): 30, 4, 9; *cockfieldensis* (Eoc.-Texas), 9(4): 95, 10, 11; *hantkeni*, var. *spissa* (Eoc.-So. Car.), 7(3): 58, 7, 13; *insecta* (Eoc.-Ala.), 8(3): 65, 8, 13, 14; *jacksonensis* (Eoc.-Ala.), 9(1): 10, 1, 23; *limbato-striata* (Plioc.-Fiji), 7(2): 30, 4, 8; *miocenica* (Mioc.-Calif.), 7(1): 8; var. *stella* (Rec.-Calif.), 6(3): 56, 7, 17; *wilcoxensis* (Eoc.-Ala.), 8(3): 65, 8, 12.

[North Carolina]. New Foraminifera from the Upper Jackson Eocene of the Southeastern Coastal Plain Region of the United States, 9(1): 1.

Nubeculinella, 9(4): 84; *bigoti* (Jurassic-France), 6(4): 84, 11, 9.

[Oligocene]. *Glandulina ozawai* Cushman, New Species, 7(4): 83. Some New Tertiary Foraminifera from Texas, 7(3): 51.

Orbitocyclina, 6(4): 92; *minima* (Cret.-Amer.), 6(4): 92.

Orbulina universa (Rec.-Calif.), 6(3): 62.

Orthovertella, 6(4): 81; *protea* (Penn.-Texas), 6(4): 81, 11, 1, 2.

Ozawaia, 7(4): 80; *tongaensis* (Rec.-Pac.), 7(4): 80, 10, 7-10.

- [Pacific]. A Recent Gümbelitria (?) from the Pacific, 10(4) : 105. Some New Recent Foraminifera from the Tropical Pacific, 9(4) : 77. Some Recent Angulogerinas from the Eastern Pacific, 8(2) : 44. The Relationships of Ungulatella, with Descriptions of Additional Species, 10(4) : 101. Two New Foraminiferal Genera from the South Pacific, 7(4) : 78.
- [Palaeozoic]. Notes on Early Palaeozoic Foraminifera, 6(2) : 43. The Species *Endothyra baileyi* (Hall), 7(4) : 90.
- Paleopolymorphina*, 6(4) : 86; *pleurostomelloides* (L. Cret.-Germany), 6(4) : 86, 12, 5.
- Parrina, A New Generic Name, 7(1) : 20.
- Parrina, 7(1) : 20.
- Patellina, Some Notes on the Genus, 6(1) : 11.
- Patellina advena* (Rec.-Pac.), 6(1) : 16, 3, 2; var. *altiformis* (Rec.-Pac.), 9(4) : 87, 9, 8; *antiqua* (Cret.-Texas), 6(1) : 12, 3, 3; *corrugata* (Rec.-Atl.), 6(1) : 15, 3, 5; *dentata* (Eoc.-France), 6(1) : 14, 3, 4; *protea* (Permian-Texas), 6(1) : 11; *subcretacea* (L. Cret.-Texas), 6(1) : 10, 3, 1; 12.
- Patellinella fijiana* (Rec.-Pac.), 9(4) : 87, 9, 7.
- Pegidia*, 6(4) : 90; *dubia* (Rec.-Pac.), 6(4) : 90, 12, 9.
- Peneroplis arietinus* (Rec.-Medit.), 7(3) : 71, 9, 19; *planatus* (Rec.-Medit.), 7(3) : 71, 9, 17, 18; *semi-lituus* (Rec.-Medit.), 7(3) : 71, 9, 20, 21.
- Pernerina depressa*, 10(2) : 36, 6, 8.
- Physalidia*, 6(4) : 88; *simplex* (Rec.-Pac.), 6(4) : 88, 12, 6.
- Planularia auris* (Rec.-Medit.), 7(3) : 70, 9, 16; *cooperensis* (Eoc.-So. Car.), 9(1) : 5, 1, 10; sp(?) (Mioc.-Calif.), 10(1) : 3, 1, 5.
- Planulina cocoaensis*, var. *cooperensis* (Eoc.-So. Car.), 9(1) : 20, 2, 12; *ornata* (Rec.-Calif.), 6(3) : 62.
- Plectina watersi* (Cret.-Texas), 9(3) : 57, 7, 1.
- Plectofrondicularia from Florida, A New, 7(3) : 60.
- Plectofrondicularia cookei* (Eoc.-So. Car.), 9(1) : 11, 1, 26; *mansfieldi* (Mioc.-Fla.), 7(3) : 60, 8, 1; *miocenica* (Mioc.-Calif.), 7(1) : 8, 1, 28.
- [Pleistocene]. Pleistocene Foraminifera from Maryland, 6(4) : 94.
- Pleurostomella austinana* (Cret.-Texas), 9(3) : 64, 7, 13; *watersi* (Cret.-Texas), 9(3) : 63, 7, 11, 12.
- [Pliocene]. New Late Tertiary Foraminifera from Vitilevu, Fiji, 7(2) : 25.
- Plummerinella*, 6(4) : 84; *complexa* (Penn.-Texas), 6(4) : 84, 11, 8.

- Polymorphina charlottensis* (Rec.-Calif.), 6(3) : 53.
Polytaxis, 6(4) : 85; *laheeii* (Penn.-Texas), 6(4) : 85, 11, 10.
Protobotellina, 6(4) : 76; *cylindrica* (Rec.-So. Atl.), 6(4) : 76, 10, 5-7.
Pseudoglandulina, 6(4) : 85; *comata* (Rec.-Medit.), 6(4) : 85, 10, 16; (Rec.-Medit.), 7(3) : 64, 8, 6; *glans* (Rec.-Medit.), 7(3) : 64, 8, 7, 8.
Pseudopolymorphina wilcoxensis (Eoc.-Ala.), 8(3) : 61, 8, 5, 6.
Pseudovigerina wilcoxensis (Eoc.-Ala.), 8(3) : 66, 8, 18; sp(?) (Cret.-Antigua), 7(2) : 40, 6, 1.
Pullenia salisburyi (Rec.-Calif.), 6(3) : 61, 8, 13; (Mioc.-Calif.), 7(1) : 15, 2, 15.
Pulvinulinella capitanensis (Mioc.-Calif.), 10(1) : 16, 3, 3; *exigua*, var. *obtusa* (Eoc.-Ala.), 8(3) : 71, 9, 9; *pulchra* (Rec.-Pac.), 9(4) : 92, 9, 10; *smithi* (Rec.-Calif.), 6(3) : 60; (Mioc.-Calif.), 7(1) : 14, 2, 12; *umbonifera* (Rec.-Pac.), 9(4) : 90, 9, 9.
Pyropilus, Notes on the Genus Tretomphalus, with Descriptions of Some New Species and a New Genus, 10(4) : 79.
Pyropilus, 10(4) : 100; **rotundatus** (Rec.-Pac.), 10(4) : 100, 13, 1, 2.
Quadrulina, 6(4) : 86; *rhabdogonoides* (Jurassic-Cret.-Europe), 6(4) : 86, 12, 4.
Quinqueloculina flexuosa (Rec.-Calif.), 6(3) : 52; (Pleistoc.-Md.), 6(4) : 95, 13, 2; *jamaicensis* (Eoc.-Jamaica), 7(4) : 76, 10, 2, 3; *lamarckiana* (Rec.-Calif.), 6(3) : 52; *seminula* (Rec.-Calif.), 6(3) : 52; (Pleistoc.-Md.), 6(4) : 95, 13, 1.
Raibosammina, 6(4) : 74; *mica* (Pal.-Okla.), 6(2) : 6, 11; (Pal.-Amer.), 6(4) : 74, 10, 3.
[Recent]. A Recent Gumbelitria (?) from the Pacific, 10(4) : 105. Additional Localities of the "Challenger" Foraminifera, 7(2) : 46. Notes on the Genus Tretomphalus, with Descriptions of Some New Species and a New Genus, Pyropilus, 10(4) : 79. Some New Recent Foraminifera from the Tropical Pacific, 9(4) : 77. Some Recent Angulogerinas from the Eastern Pacific, 8(2) : 44. Some Recent Foraminifera from off San Pedro, California, 6(3) : 49. The Relationships of Ungulatella, with Descriptions of Additional Species, 10(4) : 101. Two New Foraminiferal Genera from the South Pacific, 7(4) : 78.
Rectocibicides, 8(1) : 2; **miocenicus** (Mioc.-Fla.), 8(1) : 2, 1, 5-7.
Rectogumbelina, A New Genus from the Cretaceous, 8(1) : 4.
Rectogumbelina, 8(1) : 6; **cretacea** (Cret.-Ark.), 8(1) : 6, 1, 11, 12; **texana** (Cret.-Texas), 8(1) : 6, 1, 8-10.

- Reophax dentaliniformis* (Rec.-Calif.), 6(3) : 51, 7, 2; *excentricus* (Rec.-Calif.), 6(3) : 51; *scorpiurus* (Rec.-Calif.), 6(3) : 51, 7, 1.
- Reussia eocena* (Eoc.-Fla.), 9(1) : 13, 1, 25.
- Robertina wilcoxensis* (Eoc.-Ala.), 8(3) : 66, 8, 19.
- Robulus antipodum* (Eoc.-Calif.), 10(3) : 53, 7, 5; *arcuato-striatus*, var. *carolinianus* (Eoc.-Ala.), 9(1) : 4, 1, 9; *articulatus*, var. *texanus* (Eoc.-Calif.), 10(3) : 53, 7, 2, 3; *branneri* (Mioc.-Calif.), 10(1) : 2, 1, 4; *coloratus* (Mioc.-Calif.), 10(1) : 2, 1, 3; *gutticostatus*, var. *yazooensis* (Eoc.-Ala.), 9(1) : 4, 1, 8; *gyroscalprum* (Eoc.-Calif.), 10(3) : 53, 7, 4; *isidis* (Eoc.-Ala.), 8(3) : 52, 7, 2; *limbosus*, var. *hockleyensis* (Eoc.-Calif.), 10(3) : 52, 7, 1; *mayi* (Mioc.-Calif.), 7(1) : 2, 1, 3-5; *nikobarensis* (Mioc.-Calif.), 7(1) : 2, 1, 2; *rotulus* (Eoc.-Calif.), 10(3) : 53, 7, 6; *wilcoxensis* (Eoc.-Ala.), 8(3) : 52, 7, 3; *williamsoni* (Cret.-Antigua), 7(2) : 37, 5, 2.
- Rotalia beccarii*, var. *ornata* (Pleistoc.-Md.), 6(4) : 100, 13, 15; var. *parkinsoniana* (Pleistoc.-Md.), 6(4) : 100, 13, 14; *fimbriatula* (Cret.-Colombia), 6(3) : 69, 9, 13.
- Rugidia*, 6(4) : 90; *corticata* (Rec.-Indo-Pac.), 6(4) : 90, 12, 7.
- Rupertia* (?) *floridana* (Eoc.-Fla.), 9(1) : 21, 2, 13, 14.
- Ruttenia*, A New Name for *Bonairea* Pijpers, 1933, 9(2) : 30.
- Ruttenia*, 9(2) : 30.
- Saracenaria arcuata*, var. *hantkeni* (Eoc.-So.Car.), 9(1) : 4, 1, 11, 12; *wilcoxensis* (Eoc.-Ala.), 8(3) : 54, 7, 9.
- Schizammia*, 6(4) : 73; *labyrinthica* (Rec.-So. Atl.), 6(4) : 73, 10, 1.
- Sigmoidella plummerae* Cushman and Ozawa, The Range of, A Correction, 6(4) : 101.
- Sigmoidella plummerae* (Eoc.-Texas), 6(4) : 101.
- Sigmomorphina alabamensis* (Eoc.-Ala.), 8(3) : 62, 8, 8; *frondiculariformis* (Rec.-Calif.), 6(3) : 54; *trilocularis* (Rec.-Calif.), 6(3) : 54, 7, 16; *wilcoxensis* (Eoc.-Ala.), 8(3) : 61, 8, 7.
- Silicosigmoilina*, 6(4) : 80; *californica* (Cret.-Calif.), 6(4) : 80, 10, 17, 18.
- Sinzowella*, 9(2) : 33; *deformis*, 9(2) : 33, 3, 10-12.
- Siphogenerina mayi* (Mioc.-Calif.), 7(1) : 10, 2, 7; *nodifera* (Mioc.-Calif.), 10(1) : 13, 2, 15, 16; *pseudococcaensis* (Mioc.-Calif.), 10(1) : 13, 2, 14; *tenua* (Mioc.-Calif.), 10(1) : 13, 2, 13; *transversa* (Mioc.-Calif.), 7(1) : 10, 2, 5, 6.
- Siphogenerinoides ewaldi* (Cret.-Venezuela), 6(3) : 68, 9, 10-12.
- Siphonina carltoni* (Eoc.-La.), 8(2) : 42, 6, 5; *claibornensis* (Eoc.-Calif.), 10(3) : 62; *wilcoxensis* (Eoc.-Ala.), 8(3) : 70, 9, 7.
- Siphonodosaria fijiensis* (Plioc.-Fiji), 7(2) : 30, 4, 10.

[South Carolina]. New Foraminifera from the Upper Jackson Eocene of the Southeastern Coastal Plain Region of the United States, 9(1) : 1.

Spandelina, 6(4) : 80; *excavata* (Penn.-Texas), 6(4) : 80, 10, 14.

Spandelinoides, 6(4) : 80; *nodosariformis* (Penn.-Texas), 6(4) : 80, 10, 13.

Sphaeridia, 6(4) : 90; *papillata* (Rec.-Indo-Pac.), 6(4) : 90, 12, 8.

Spiroplectammina anceps, 8(4) : 89, 90, 11, 5, 6; *baudouiniana*, 8(4) : 87, 90, 11, 1; *dentata*, 8(4) : 91, 11, 7; *excolata*, 8(4) : 94; *laevis*, 8(4) : 87, 11, 2; var. *cretosa* (Cret.-Texas), 8(4) : 87, 11, 3; *navarroana* (Cret.-Texas), 8(4) : 96, 11, 14; *obtusangula*, 8(4) : 88, 11, 4; *parallela* (Plioc.-Fiji), 7(2) : 26, 4, 1;

scotti (L. Cret.-Texas), 6(1) : 8, 2, 11; *semicomplanata*, 8(4) : 94, 11, 8, 9; var. *junceae* (Cret.-Ark.), 8(4) : 95, 11, 10; *whitneyi*

(L. Cret.-Texas), 6(1) : 8, 2, 12; *wilcoxensis* (Eoc.-Ala.), 8(3) : 51, 7, 1.

Spiroplectoides and Its Species, Notes on the Genus, 10(2) : 37.

Spiroplectoides from the Cretaceous of California, A New, 10(3) : 70.

Spiroplectoides, 10(2) : 37; *californica*, 10(3) : 70, 9, 15-17;

clotho, 10(2) : 42, 6, 19-23; *curta* (Eoc.-So. Car.), 9(1) : 11, 1,

24; 10(2) : 43, 6, 27; *eocenica*, 10(2) : 43, 6, 28; *flexuosa*, 8(4) : 91; 10(2) : 41, 6, 14-18; *papillata*, 10(2) : 41, 6, 24; *rosula*,

10(2) : 38, 6, 10-13; *spectabilis*, 10(2) : 43, 6, 25, 26.

Stegnammina, 6(4) : 74; *cylindrica* (Pal.-Amer.), 6(4) : 74, 10, 2.

[Structural]. Notes on Early Palaeozoic Foraminifera, 6(2) : 43.

The Development of Hantkenina in the Cretaceous with a Description of a New Species, 6(2) : 39. The Species *Endothyra*

baileyi (Hall), 7(4) : 90.

[Taxonomic]. A Resumé of New Genera of the Foraminifera

Erected Since Early 1928, 6(4) : 73. *Dentostomina*, A New

Genus of the Miliolidae, 9(2) : 31. Notes on Some of the

Earlier Species Originally Described as *Bulimina*, 10(2) : 27.

Notes on the Foraminifera Described by Batsch in 1791, 7(3) :

62. Notes on the Genus *Spiroplectoides* and Its Species, 10(2) :

37. Notes on the Genus *Tretomphalus*, with Descriptions of

Some New Species and a New Genus, *Pyropilus*, 10(4) : 79.

Notes on the Genus *Virgulina*, 8(1) : 7. On Homonyms in

Foraminifera, 9(4) : 96. On *Uvigerina pigmea* d'Orbigny,

6(3) : 62. *Parrina*, A New Generic Name, 7(1) : 20. Relations

and Geologic Distribution of the Genera of the *Valvulinidae*,

9(2) : 38. *Ruttenia*, A New Name for *Bonairea Pijpers*,

1933, 9(2) : 30. Some New Foraminiferal Genera, 9(2) : 32.

Some Notes on d'Orbigny's Models, 9(3) : 70. Some Notes on the Genus Flabellinella Schubert, 9(1) : 16. Some Notes on the Genus Patellina, 6(1) : 11. Textularia and Related Forms from the Cretaceous, 8(4) : 86. The Generic Position of "Cornuspira cretacea Reuss," 10(2) : 44. The Genus Vulvulina and Its Species, 8(4) : 75. The Microspheric and Megalospheric Forms of Valvulina oviedoina d'Orbigny, 7(1) : 17. The Relationships of Textulariella and Description of a New Species, 8(4) : 97. The Relationships of Ungulatella, with Descriptions of Additional Species, 10(4) : 101. The Species Endothyra baileyi (Hall), 7(4) : 90.

[Tennessee]. New American Cretaceous Foraminifera, 9(3) : 49.

[Texas]. A New Species of Clavulina from the Cretaceous of Texas, 9(1) : 21. Additional New Eocene Foraminifera, 8(2) : 40. Hastigerinella and Other Interesting Foraminifera from the Upper Cretaceous of Texas, 7(4) : 83. New American Cretaceous Foraminifera, 9(3) : 49. Notes on Upper Cretaceous Species of Vaginulina, Flabellina and Frondicularia from Texas and Arkansas, 6(2) : 25. Rectogümbelina, A New Genus from the Cretaceous, 8(1) : 4. Some New Tertiary Foraminifera from Texas, 7(3) : 51. Some Vaginulinas and Other Foraminifera from the Lower Cretaceous of Texas, 6(1) : 1. Two New Navarro Foraminifera from Texas, 8(4) : 98. Two New Texas Foraminifera, 9(4) : 95.

Textularia and Related Forms from the Cretaceous, 8(4) : 86.

Textularia brevicona, 8(4) : 92; *decurrens*, 8(4) : 92; *ehrenbergii*, 8(4) : 87; *laevigata* (Mioc.-Calif.), 8(2) : 32, 5, 6, 7; *mexicana* (Mioc.-Calif.), 8(2) : 33, 5, 3; *mississippiensis*, var. *rhomboidea* (Eoc.-Texas), 7(3) : 51, 7, 1; *nactaensis*, 8(4) : 96; *obsoleta*, 8(4) : 90; *parallela*, 8(4) : 94; *ripleyensis*, 8(4) : 96, 11, 12, 13; *sagittula*, var. *coonensis*, 8(4) : 96; *schencki* (Rec.-Calif.), 6(3) : 51; *serrata*, 8(4) : 91; *smithvillensis* (Eoc.-Texas), 9(4) : 95, 10, 10; *subconica*, 8(4) : 95; 11, 11; *subglabra*, 8(4) : 94; *triquetra*, 8(4) : 89; *trochus*, 8(4) : 86; *turris*, 8(4) : 87; *warreni* (Olig.-Texas), 7(3) : 51, 7, 2.

Textulariella and Description of a New Species, The Relationships of, 8(4) : 97.

Textulariella cretosa (Cret.-England), 8(4) : 97, 11, 17-19.

Thurammina irregularis (Pal.-Okla.), 6(2) : 6, 8; *phasela* (Pal.-Okla.), 6(2) : 6, 7.

Trepeilopsis, 6(4) : 76; *grandis* (Penn.-Texas), 6(4) : 76, 10, 10.

Tretomphalus, with Descriptions of Some New Species and a

- New Genus, *Pyropilus*, Notes on the Genus, 10(4) : 79.
- Tretomphalus atlanticus* (Rec.-Atl.), 10(4) : 86, 11, 3; 12, 7; *bulloides* (Rec.-Atl.), 10(4) : 86, 11, 1, 2; 12, 6; *clarus* (Rec.-Pac.), 10(4) : 99, 11, 6; 12, 16, 17; *concinus* (Rec.-Pac.), 10(4) : 96, 11, 8, 9; 12, 13-15; *grandis* (Rec.-Pac.), 10(4) : 95, 11, 10; 12, 23, 24; *milletti* (Rec.-Pac.), 10(4) : 87, 11, 4, 5; 12, 1-5; *pacificus* (Rec.-Pac.), 10(4) : 93, 11, 7; 12, 8-12; *planus* (Rec.-Pac.), 10(4) : 94, 11, 11; 12, 18-22.
- Triloculina from the Claiborne of Louisiana, New Species of, 10(3) : 65.
- Triloculina inusitata* (Eoc.-La.), 10(3) : 66, 9, 8-10; *mirifica* (Eoc.-La.), 10(3) : 65, 9, 11-14; *oblonga* (Rec.-Calif.), 6(3) : 52; *paulocostata* (Eoc.-La.), 10(3) : 69, 9, 6, 7; *rotunda* (Pleistoc.-Md.), 6(4) : 95, 13, 3.
- Trimosina orientalis* (Rec.-Pac.), 9(4) : 78, 8, 4.
- [Trinidad]. Fossil Species of *Hastigerinella*, 6(1) : 17. Post-Cretaceous Occurrence of *Gümbelina* with a Description of a New Species, 9(3) : 64. Some Interesting New Uniserial Foraminifera from Trinidad, 10(3) : 71.
- Triplasia wilcoxensis* (Eoc.-Ala.), 8(3) : 58, 7, 19.
- Tritaxia tricarinata*, 8(4) : 89.
- Trochammina globigeriniformis* (Rec.-Calif.), 6(3) : 52, 7, 11; *teasi* (Eoc.-Texas), 7(3) : 52, 7, 3; *turbinata* (Rec.-Calif.), 6(3) : 53, 7, 10.
- Tuberitina*, 6(4) : 93; *bulbacea* (Penn.-Okla., Texas), 6(4) : 93, 12, 13, 14.
- Tubulogenerina eocenica* (Eoc.-Texas), 8(2) : 42, 6, 4.
- Ungulatella, with Descriptions of Additional Species, The Relationships of, 10(4) : 101.
- Ungulatella, 7(4) : 81; 10(4) : 102; *capistra* (Rec.-Pac.), 10(4) : 103, 13, 8; *conoides* (Rec.-Pac.), 10(4) : 102, 13, 6, 7; *pacifica* (Rec.-Pac.), 7(4) : 82, 10, 11, 12; (Rec.-Pac.), 10(4) : 102, 13, 3, 4; *peregrina* (Rec.-Pac.), 10(4) : 102, 13, 5.
- Uvigerina pigmea d'Orbigny, On, 6(3) : 62.
- Uvigerina carmelensis* (Mioc.-Calif.), 10(1) : 11, 2, 7; *glabrans* (Eoc.-Ala.), 9(1) : 13, 1, 28; *hootsi* (Mioc.-Calif.), 10(1) : 22, 3, 8, 9; *modeloensis* (Mioc.-Calif.), 10(1) : 12, 2, 8; *pigmea* (Plioc.-Italy), 6(3) : 62, 9, 14-20; var. *curticostata* (Rec.-Calif.), 6(3) : 60; *proboscidea* (Mioc.-Calif.), 10(1) : 12, 2, 12; var. *vadescens* (Rec.-Pac.), 9(4) : 85, 8, 14, 15; *segundoensis* (Mioc.-Calif.), 10(1) : 26, 4, 11; *senticosa* (Rec.-Calif.), 6(3) : 58; *subperegrina* (Mioc.-Calif.), 10(1) : 12, 2, 9-11; *vicksburg-*

ensis (Olig.-Texas), 7(3): 54, 7, 7; *yazooensis* (Eoc.-Ala.), 9(1): 13, 1, 29.

Uvigerinella californica (Mioc.-Calif.), 7(1): 10, 2, 3; var. *gracilis* (Mioc.-Calif.), 10(1): 11, 2, 6; *obesa* (Mioc.-Calif.), 7(1): 10, 2, 4.

Vaginulina, Flabellina and Frondicularia from Texas and Arkansas, Notes on Upper Cretaceous Species of, 6(2): 25.

Vaginulina costifera (Eoc.-Calif.), 10(3): 54, 7, 9; *intumescens* (L. Cret.-Texas), 6(1): 5, 2, 1-6; *kochii* (L. Cret.-Texas), 6(1): 1, 1, 1-9; var. *striolata* (L. Cret.-Texas), 6(1): 4, 1, 10-16; *leguminiformis* (Rec.-Medit.), 7(3): 68, 8, 24; *margaritifera* (Rec.-Medit.), 7(3): 69, 9, 7-9; *marginulinoides* (L. Cret.-Texas), 6(1): 5, 1, 23-26; *multicostata* (Cret.-Ala.), 6(2): 28, 4, 4; *recta* (Cret.-Texas), 6(2): 27, 4, 5; (L. Cret.-Texas), 6(1): 4, 1, 17-22; *simondsi* (Cret.-Ark.), 6(2): 27, 4, 6-8; *strigillata* (Cret.-Texas, Ark.), 6(2): 26, 4, 1, 9, 10; *texana* (Cret.-Texas), 6(2): 30, 4, 2, 3; *trilobata* (Cret.-Texas), 6(2): 30, 4, 11; *webbervillensis* (Cret.-Texas, Ark.), 6(2): 27, 4, 14; *wrighti* (Eoc.-Calif.), 10(3): 54, 7, 8; sp(?) (Cret.-Texas), 6(2): 30, 4, 12, 13.

Vaginulinas and Other Foraminifera from the Lower Cretaceous of Texas, Some, 6(1): 1.

Valvulammina, 9(2): 37; *globulosa*, 9(2): 37, 4, 14-16.

Valvulina oviedoiana d'Orbigny, The Microspheric and Megalospheric Forms of, 7(1): 17.

Valvulina oviedoiana (Rec.-Atl.), 7(1): 17, 3, 6-10.

Valvulineria allomorphinoides (Cret.-Antigua), 7(2): 43, 6, 2; *araucana* (Rec.-Calif.), 6(3): 60, 8, 10; *grandis* (Mioc.-Calif.), 10(1): 26, 4, 12; *involuta* (Eoc.-Calif.), 10(3): 63, 8, 12; *jacksonensis* (Eoc.-Ala.), 9(1): 18, 2, 9; *joaquinensis* (Mioc.-Calif.), 10(1): 14, 2, 17; *miocenica*, var. *depressa* (Mioc.-Calif.), 7(1): 11, 2, 8; *scrobiculata* (Eoc.-Ala.), 8(3): 70, 9, 5; *texana* (Eoc.-Texas), 7(3): 56, 7, 9; *vilardeboana*, var. *glabra* (Rec.-Calif.), 6(3): 60, 8, 9; *wilcoxensis* (Eoc.-Ala.), 8(3): 70, 9, 6.

[Venezuela]. Notes on Some Foraminifera from Venezuela and Colombia, 6(3): 64.

Verneuilina advena (Rec.-Calif.), 6(3): 52, 7, 6; *cretosa* (Cret.-Texas), 9(3): 51, 5, 7; *polystropha*, 10(2): 30, 5, 17; *schizea* (L. Cret.-Texas), 6(1): 9, 2, 13, 14.

Vertebralina jamaicensis (Eoc.-Jamaica), 7(4): 77, 10, 5, 6.

Victoriella, 6(4): 92; *pectae* (Tert.-Australia), 6(4): 92, 12, 11.

Virgulina from the Miocene of Florida, A New, 7(2): 32. Notes on the Genus Virgulina, 8(1): 7.

Virgulina, 8(1) : 8; *acuta* (Mioc.-Europe), 8(1) : 16, 3, 4; *advena*, 8(1) : 14, 2, 11; *bradyi* (Rec.-Atl.), 8(1) : 14, 2, 12; *bramletti* (Rec.-Calif.), 6(3) : 57, 8, 2; 8(1) : 20, 3, 9; *californiensis* (Mioc.-Calif.), 8(1) : 20, 3, 8; var. *grandis* (Mioc.-Calif.), 10(1) : 9, 1, 15, 16; var. *ticensis* (Mioc.-Calif.), 10(1) : 10, 1, 17; *carinata*, 8(1) : 14, 2, 10; *compressa* (Rec.-Atl.), 8(1) : 10, 2, 2; *cornuta* (Rec.-Calif.), 6(3) : 57, 8, 1; 8(1) : 14, 2, 9; *delmonteensis* (Mioc.-Calif.), 10(1) : 25, 4, 9; *dibollensis* (Eoc.-U. S.), 8(1) : 21, 3, 14; *digitalis*, 8(1) : 21, 3, 12; *fusiformis* (Mioc.-Fla.), 8(1) : 20, 3, 10; *gunteri*, var. *curtata* (Mioc.-Fla.), 8(1) : 4, 1, 2; *hungarica* (Eoc.-Hungary), 8(1) : 21; *mexicana* (Rec.-Atl.), 8(1) : 15, 2, 13; *miocenica* (Mioc.-Fla.), 7(2) : 32, 4, 14-16; *mustoni* (Olig.-Europe), 8(1) : 20, 3, 11; *navarroana* (Cret.-Texas), 9(3) : 63, 7, 9, 10; *nodosa* (Plioc.-Calif.), 8(1) : 16, 3, 3; *obscura*, 8(1) : 11, 2, 7; *pauciloculata*, 8(1) : 10, 2, 3; *pontoni* (Mioc.-Fla., Calif.), 8(1) : 17, 3, 7; *punctata* (Rec.-Mioc.), 8(1) : 9, 2, 1; *recta* (Eoc.-So. Car.), 9(1) : 12, 1, 31; *schreibersiana*, 8(1) : 17, 3, 5; var. *complanata*, 8(1) : 11, 2, 6; var. *marginata* (Eoc.-Nigeria), 8(1) : 21, 3, 13; var. *obesa* (Plioc.-England), 8(1) : 16, 3, 2; *squammosa*, 8(1) : 15, 3, 1; *subdepressa*, 8(1) : 10, 2, 4; *subsquamosa* (Mioc.-Bavaria), 8(1) : 17, 3, 6; var. *striata*, 8(1) : 12, 2, 8; *tenuis* (Pleistoc.-Sicily), 8(1) : 15, 2, 14; *textulata*, 8(1) : 11, 2, 5; *wilcoxensis* (Eoc.-Ala.), 8(3) : 67, 8, 22; *zetina* (Eoc.-Mexico), 8(1) : 22, 3, 15.

Virgulina (*Virgulinella*) *gunteri* (Mioc.-Fla.), 8(1) : 22, 3, 17; var. *curtata* (Mioc.-Fla.), 8(1) : 22, 3, 18; *miocenica* (Mioc.-Fla.), 8(1) : 23, 3, 19; *pertusa*, 8(1) : 22, 3, 16.

Virgulinella, 8(1) : 9.

Vulvulina and Its Species, The Genus, 8(4) : 75.

Vulvulina, 8(4) : 75; *advena* (Eoc.-Ala.), 8(4) : 84, 10, 14; *arenacea* (Rec.-Pac.), 8(4) : 79, 10, 13; *colei* (Eoc.-Mexico), 8(4) : 84, 10, 21, 22; *flabelliformis* (Eoc.-Europe), 8(4) : 83, 10, 18, 19; *jarvisi* (Eoc.-Trinidad), 8(4) : 84, 10, 20; *nummulina* 8(4) : 82, 10, 16; *pacifica* (Rec.-Pac.), 8(4) : 78, 10, 8, 9; *pectinata*, 8(4) : 82, 10, 12; var. *mexicana* (Eoc.-Mexico), 8(4) : 83, 10, 17; *pennatula* (Rec.-Medit.), 7(3) : 69, 9, 10-13; 8(4) : 76, 10, 1-5; var. *italica* (Mioc.-Italy), 8(4) : 78, 10, 6, 7; *spinosa* (Eoc.-Amer.), 8(4) : 79, 10, 15; var. *miocenica* (Mioc.-Amer.), 8(4) : 80, 10, 10.

Wiesnerella, 9(2) : 33; *auriculata*, 9(2) : 33, 3, 7-9.

Yaberinella, 6(4) : 77; *jamaicensis* (Eoc.-Jamaica), 6(4) : 77.

